

Is Nöther Symmetric Approach Consistent With Dynamical Equation In Non-Minimal Scalar-Tensor Theories?

A.K.Sanyal^{*,1}, B.Modak^{**,2}

February 7, 2008

¹ Dept. of Physics, Jangipur College, Murshidabad, West Bengal, India - 742213.

² Dept. of Physics, University of Kalyani,
India - 741235.

1

Abstract

The form of the coupling of the scalar field with gravity and the potential have been found by applying Nöther theorem to 2-dimensional minisuperspaces in induced gravity model. It has been observed that though the forms thus obtained are consistent with all the equations $\mathcal{L}_X L = 0$, yet they do not satisfy the field equations for $k = \pm 1$, in Robertson-Walker model. It has been pointed out that this is not due to the degeneracy of the Lagrangian, since this problem does not appear in $k = 0$ case. It has also been shown that though Nöther theorem fails to extract any symmetry from the Lagrangian of such model for $k = \pm 1$, symmetry exists, which can easily be found by studying the continuity equation.

Keywords : Induced gravity theory, Nöther symmetry ; cosmology.

1 Introduction

The idea of applying Nöther theorem to cosmological minisuperspaces is not far too old. In 1990, De Ritis et al [1], perhaps for the first time applied it to the homogeneous isotropic cosmological models of general theory of relativity with a minimally coupled scalar field. Since then, lot of works came out in the literature [2]. During early nineties, theory of gravitation with nonminimally coupled scalar field was supposed to encounter a number of longstanding puzzles of cosmology, e.g., it was thought to produce successful phase transition for a large class of models and help to solve the graceful exit problem on one hand and density perturbation on the other [3].

However, the problem with such model is that, the exact form of the coupling of the scalar field with gravity is not known a priori. Capozziello et al [4] following [1] tried to find the form of the coupling $f(\phi)$ along with the potential $V(\phi)$ for a scalar field ϕ by applying Nöther theorem in induced gravity theory. Nöther theorem states that, if there exists a vector field X , for which the Lie derivative of a given Lagrangian L , vanishes ($\mathcal{L}_X L = 0$), the Lagrangian admits Nöther symmetry yielding a conserved current. The idea of the theorem under the present context is to find the form of $f(\phi)$ and $V(\phi)$ for which there would exist a vector field X that would make $\mathcal{L}_X L = 0$ and hence would yield a conserved current. It is not expected in general that, in all such models there would exist certain $f(\phi)$ and $V(\phi)$ and hence X , that would make $\mathcal{L}_X L = 0$. Rather, while solving the set of equations $\mathcal{L}_X L = 0$, one might some time encounter inconsistencies which would imply that the Lagrangian does not admit Nöther symmetry in such models. However, it is generally believed that if certain $f(\phi)$ and $V(\phi)$ and hence X exists, that satisfy all the equations $\mathcal{L}_X L = 0$, then the Lagrangian admits Nöther symmetry.

The last statement is not found to be true in general. In the following section we shall review the work of Capozziello et al [4] and show, what they might have overlooked, that for induced theory of gravity in the Robertson-walker minisuperspace model, there apparently exists, Nöther symmetry in the sense that all the

¹Relativity and Cosmology Research Centre
Department of Physics, Jadavpur University
Calcutta - 700 032, India
Electronic address: * aks@juphys.ernet.in,
** bmodak@klyuniv.ernet.in

equations $\mathcal{L}_X L = 0$ are satisfied, together with the existence of a conserved current. However, the functions $f(\phi)$ and $V(\phi)$ thus obtained are not found to satisfy the field equations for $k = \pm 1$. This result perhaps is unique and not been encountered earlier. It is true, as pointed out by Capozziello et al [4], that the form of $f(\phi)$ makes the Lagrangian degenerate. However, it is known that a degenerate Lagrangian gives rise to a constrained system for which Legendre transformation does not exist and Hamiltonian dynamics can not be produced unless the constraint is tackled properly [5]. Further, as far as Lagrangian dynamics is concerned, degeneracy introduces underdeterminacy, i.e., the number of field equations are less than the number of true degrees of freedom. So degeneracy is not responsible for the inconsistency mentioned above.

We shall further show in this section that for $k = \pm 1$, apparently Nöther symmetry is found to exist with zero potential but real solution exists only for $k = -1$. Due to the degeneracy of the Lagrangian, the field equations reduce to one with two field variables in the case of $k = -1$. This equation is found to admit inflationary solution.

At the end of section 2, we shall further show that the same form of $f(\phi)$ with a different $V(\phi)$ satisfy the field equations for $k = 0$, for which all the field equations reduce to one only, viz., the constant of motion with two degrees of freedom $a(t)$ and $\phi(t)$. We shall also show that there exists nontrivial solution of $a(t)$, if certain form of $\phi(t)$ is assumed. Hence we conclude that the trivial solution $a(t) = 0, \dot{\phi}(t) = \phi(t) = 0$, obtained by Capozziello et al [4] is not due to the presence of degeneracy rather due to the use of a typical form of $f(\phi)$ and $V(\phi)$, which is the outcome of Nöther symmetry, but that do not satisfy the field equations.

The form of $f(\phi)$ we are talking of so far is only a particular solution of a first order non-linear differential equation of f that can not be solved in general, to obtain f in closed form. Since we are only interested in those Lagrangian in which gravitational coupling is in closed form, so at this stage we can always make the comment that, Nöther symmetry does not exist for such models with $k = \pm 1$, for potentials other than zero. For zero potential the symmetry is found to exist only for $k = -1$, which is consistent with the dynamical equations.

In section 3 we have chosen a particular form of $f(\phi)$, viz., $f(\phi) = \epsilon\phi^2$ keeping a parameter ϵ arbitrary and have shown that the study of the field equation viz., the continuity equation, reveals the existence of certain symmetry in Lagrangian for which there is a constant of motion and which can not be found by studying Nöther theorem. We had a belief that all dynamical symmetries can be encountered by applying Nöther theorem. However following the above result such belief is now questionable. Section 4 concludes our present work.

2 Nöther Symmetry In Robertson-Walker Model

We start with the following action,

$$A = \int d^4x (\sqrt{-g} [f(\phi)R - \frac{1}{2}(\phi_{,\mu}\phi^{,\mu} - V(\phi))]) \quad (1)$$

which for Robertson-Walker metric, reduces to

$$A = \int [-fa\dot{a}^2 - f'a^2\dot{\phi} + kaf + \frac{1}{12}a^3\dot{\phi}^2 - \frac{1}{6}a^3V]dt + \text{surface-term} \quad (2)$$

Field equations along with the Hamiltonian constraint equations are

$$2\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + 2\frac{f'a}{fa}\dot{\phi} + \frac{1}{f}[f'' + \frac{1}{4}]\dot{\phi}^2 + \frac{f'}{f}\ddot{\phi} + \frac{k}{a^2} - \frac{V}{2f} \quad (3)$$

$$\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} - \frac{\dot{a}\dot{\phi}}{2f'a} - \frac{\ddot{\phi}}{6f'} + \frac{k}{a^2} - \frac{V'}{6f'} = 0 \quad (4)$$

$$\frac{\dot{a}^2}{a^2} + \frac{f'\dot{a}\dot{\phi}}{fa} - \frac{\dot{\phi}^2}{12f} + \frac{k}{a^2} - \frac{V}{6f} = 0 \quad (5)$$

In the above, $k = 0, \pm 1$ and dot represents derivative with respect to proper time, while dash, derivative with respect to ϕ . In the Lagrangian under consideration the configuration space is $Q = (a, \phi)$, whose tangent space $TQ = (a, \phi, \dot{a}, \dot{\phi})$. Hence the infinitesimal generator of the Nöther symmetry is

$$X = \alpha \frac{\partial}{\partial a} + \beta \frac{\partial}{\partial \phi} + \dot{\alpha} \frac{\partial}{\partial \dot{a}} + \dot{\beta} \frac{\partial}{\partial \dot{\phi}} \quad (6)$$

The existence of Nöther symmetry implies the existence of the vector field X such that the Lie derivative of the Lagrangian with respect to the vector field vanishes i.e.

$$\mathcal{L}_X L = 0 \quad (7)$$

This gives an expression of second degree in a and ϕ , whose co-efficient are functions of a and ϕ only. Thus to satisfy equation (7), we have

$$\alpha f + a\beta f' + 2af \frac{\partial \alpha}{\partial a} + a^2 f' \frac{\partial \beta}{\partial a} = 0, \quad (8)$$

$$\alpha - 4\alpha' f' + \frac{2}{3}a\beta' = 0, \quad (9)$$

$$a\beta f'' + [a \frac{\partial \alpha}{\partial a} + 2\alpha + a\beta']f' - \frac{a^2}{6} \frac{\partial \beta}{\partial a} + 2\alpha' f = 0, \quad (10)$$

$$k(\alpha f + a\beta f') - \frac{a^2}{2}[\alpha V + \frac{a\beta V'}{3}] = 0. \quad (11)$$

The above set of equations can be solved by the method of separation of variables. Essentially for $k = \pm 1$, one ends up with a non linear differential equation of f which is

$$3f'^2 + f = Qf^3 \quad (12)$$

where Q is a constant of integration. For $Q = 0$, this equation can immediately be solved to yield

$$f = -\frac{1}{12}(\phi - \phi_0)^2 \quad (13)$$

α, β and V can also be obtained as a by product viz.,

$$\alpha = \frac{2n}{a(\phi - \phi_0)^2}; \beta = \frac{n}{a^2(\phi - \phi_0)}; V = \lambda(\phi - \phi_0)^6 \quad (14)$$

Since, solutions (13) and (14) satisfy the set of equations (8) - (11) i.e. equation (7), so one can now claim that the Lagrangian (2) admits Nöther symmetry for the above form of f and V . Now for the Cartan one form

$$\theta_L = \frac{\partial L}{\partial a} da + \frac{\partial L}{\partial \phi} d\phi, \quad (15)$$

the constant of motion $i_X \theta_L$ is obtained as,

$$F = \frac{\frac{d}{dt}[a(\phi - \phi_0)]}{\phi - \phi_0}. \quad (16)$$

So far, we have produced nothing new. However, at this stage Capozziello et al [4] went on to find cyclic co-ordinate and have produced a trivial solution like $a(t) = 0 = \phi(t)$. It is true that the form of f given by (13), makes the Lagrangian (2) degenerate, since its Hessian determinant vanishes. Degenerate Lagrangian leads to constrained dynamics for which Legendre transformation does not exist and hence Hamiltonian can not be expressed. In the context of general relativity degenerate Lagrangian appears quite often leading to underdetermined situation, i.e. degrees of freedom becomes more in number than the field equations and relations like equation of state etc. are assumed to tackle the situation. However, in the following we shall analyse the constraint to inspect the outcome. For this purpose we find f', f'' and V' from equations (13) and (14) and substitute all these together with f and V in equations (3), (4) and (5) to obtain.

$$2\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + 4\frac{\dot{a}\dot{\phi}}{a(\phi - \phi_0)} - \frac{\dot{\phi}^2}{(\phi - \phi_0)^2} + 2\frac{\ddot{\phi}}{\phi - \phi_0} + \frac{k}{a^2} + 6\lambda(\phi - \phi_0)^4 = 0, \quad (17)$$

$$\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + 3\frac{\dot{a}\dot{\phi}}{a(\phi - \phi_0)} + \frac{\ddot{\phi}}{\phi - \phi_0} + \frac{k}{a^2} + 6\lambda(\phi - \phi_0)^4 = 0, \quad (18)$$

$$\frac{\dot{a}^2}{a^2} + 2\frac{\dot{a}\dot{\phi}}{a(\phi - \phi_0)} + \frac{\dot{\phi}^2}{(\phi - \phi_0)^2} + \frac{k}{a^2} + 2\lambda(\phi - \phi_0)^4 = 0. \quad (19)$$

Now, taking the time derivative of equation (16) and using it to eliminate $\ddot{a}$ and $\ddot{\phi}$ terms from equation (17) or (18) one ends up with the following equation in either case, viz.,

$$\frac{\dot{a}^2}{a^2} + 2\frac{\dot{a}\dot{\phi}}{a(\phi - \phi_0)} + \frac{\dot{\phi}^2}{(\phi - \phi_0)^2} + \frac{k}{a^2} + 6\lambda(\phi - \phi_0)^4 = 0. \quad (20)$$

It is now evident that equations (19) and (20) lead to inconsistency unless $\lambda = 0$. This inconsistency can also be viewed other way round. Eliminating $\ddot{a}$ and k/a^2 terms from equations (3), (4) and (5), which is achieved by multiplying equation (4) by 2, subtracting it from equation (3) and finally adding it up with equation (5), one obtains

$$(3f'^2 + f)[\ddot{\phi} + 3\frac{\dot{a}\dot{\phi}}{a}] + (3f'^2 + f)'\frac{\dot{\phi}^2}{2} + (fV' - 2Vf') = 0. \quad (21)$$

This is an important relation that we shall come across in the following section to find symmetries other than Nöther symmetry. However, presently we observe that for $3f'^2 + f = 0$, which is the equation (12) for $Q = 0$ and in particular for solution (13) of f , the potential is in the form

$$V = \lambda(\phi - \phi_0)^4, \quad (22)$$

λ being a constant of integration. It is now apparent that for the solution (13) of f , field equations are satisfied for a potential given by the form (22) and not by the form (14). It can further be shown that the potential given by the form (22) does not yield conserved current, which implies Nöther symmetry does not exist for this case. To show this, we construct yet another equation by subtracting equation (4) from equation (3) viz.,

$$\frac{\ddot{a}}{a} + [2\frac{f'}{f} + \frac{1}{2f'}]\frac{\dot{a}}{a}\dot{\phi} + [\frac{f'}{f} + \frac{1}{4f}]\dot{\phi}^2 + [\frac{f'}{f} + \frac{1}{6f'}]\ddot{\phi} + \frac{V'}{6f'} - \frac{V}{2f} = 0. \quad (23)$$

For the form of f and V given by (13) and (14), one obtains the conserved current given by (16). However for any form of the potential other than zero, no such conserved current can be obtained, confirming our claim stated above. At this point we like to mention that we have come across a very important result that has never been encountered earlier and that is, even if a Lagrangian can be found to admit Nöther symmetry, it might not satisfy the field equations. Truly, in view of equations (19) and (20) we can say that it does not satisfy the Hamiltonian Constraint equation in particular, which is a typical feature of gravitation, and hence the result remained unnoticed. We further like to mention that the trivial solutions obtained by Capozziello et al [4] is not due to the degeneracy of the Lagrangian, rather due to a wrong choice of potential given by equation (14) that does not satisfy the field equations.

We have already noticed that the conserved current (16) and hence Nöther symmetry exists for $V = 0$, which further satisfies the field equations. So let us study the case. For $V = 0$, the derivative of equation (16) along with the Hamiltonian Constraint equation (19) trivially satisfy equations (17) and (18). Hence we are left with equation (16) and equation (19), with $\lambda = 0$. Now, squaring equation (16) and substituting it in equation (19) with $\lambda = 0$, one gets

$$F^2 + k = 0 \quad (24)$$

i.e. F is determined by the curvature parameter, which has to be $k = -1$ for real solution. Hence, we have come across yet another contradiction that though Nöther symmetry apparently has been found to exist for $k = \pm 1$, but for real solutions only $k = -1$ is allowed. However, even for $k = -1$, we have got only one equation viz. (16) to solve for a pair of field variables a and ϕ . This is the out come of a degenerate Lagrangian as mentioned earlier. In any case, equation (16) can be solved in principle by assuming a form of a or ϕ . To see whether equation (16) admits inflation, let us choose $a = Ae^{mt}$, A and m being constants. The solution for ϕ thus turns out to be

$$\phi = C \exp[-mt - \frac{F}{Am} \exp(-mt)], \quad (25)$$

where C is the constant of integration and ϕ_0 has been assumed to be zero. As $t \rightarrow 0$, $\phi \rightarrow ce^{-F/Am}$ and hence becomes constant while as $t \rightarrow \infty$, $\phi \rightarrow 0$. Thus the solution of ϕ is found to be well behaved.

We have already shown that for the form of f and V given by relations (13) and (22) Nöther symmetry does not exist, hence this case can be studied to show once again that degeneracy produces underdetermined equations and nothing else. In view of relations (13) and (22), field equations (3), (4) and (5) reduces to

$$2\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{\dot{a}\dot{\phi}}{a\phi} - \frac{\dot{\phi}^2}{\phi^2} + 2\frac{\ddot{\phi}}{\phi} + \frac{k}{a^2} + 6\lambda\phi^2 = 0. \quad (26)$$

$$\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + 3\frac{\dot{a}\dot{\phi}}{a\phi} - \frac{\ddot{\phi}}{\phi} + \frac{k}{a^2} + 4\lambda\phi^2 = 0. \quad (27)$$

$$\frac{\dot{a}^2}{a^2} + 2\frac{\dot{a}\dot{\phi}}{a\phi} + \frac{\dot{\phi}^2}{\phi^2} + \frac{k}{a^2} + 2\lambda\phi^2 = 0. \quad (28)$$

where, for the sake of simplicity we have chosen $\phi_0 = 0$. We further restrict ourself to the case $k = 0$. Now multiplying equation (28) by 3 and subtracting it from equation (26) one obtains,

$$\frac{\ddot{a}}{a} - \frac{\dot{a}^2}{a^2} + \frac{\ddot{\phi}}{\phi} - 2\frac{\dot{\phi}^2}{\phi^2} - \frac{\dot{a}\dot{\phi}}{a\phi} = 0, \quad (29)$$

which further yields

$$\frac{\dot{a}}{a} + \frac{\dot{\phi}}{\phi} = m\phi, \quad (30)$$

where, m is the constant of integration. Using equation (26) in equations (22), (23) and (24) one obtains only a single outcome, which is a restriction of m , viz.,

$$m^2 + 2\lambda = 0. \quad (31)$$

So m is determined. Equation (31) implies that for real solution λ should be negative. Hence we end up with a single equation viz., (30). This is what we have claimed earlier, that degeneracy in Lagrangian produces an underdetermined situation, viz., one equation for two degrees of freedom, in the present context.

One can now try to solve equation (30) for the scale factor by choosing a solution for ϕ with the requirement that the early universe was dominated by the scalar field while it vanishes asymptotically. For such requirement one can choose ϕ in the following form

$$\phi = e^{-p^2 t}, \quad (32)$$

where p^2 is a constant. Under such assumption given by equation (32) the scale factor takes the following form

$$a = l e^{[p^2 t - \frac{m}{p^2} e^{-p^2 t}]}, \quad (33)$$

where l is yet another constant of integration. The above solution shows that the scale factor vanishes as $t \rightarrow -\infty$, while it has got a constant value at $t = 0$ and exponentially increases with t .

3 No Other Symmetry In Robertson-Walker Model?

In the last section we have observed that for $k = \pm 1$, the Nöther symmetry that was supposed to exist, actually does not, since the required forms of f and V for the existence of Nöther symmetry do not satisfy the field equations except for a very special case viz., $V = 0$ and $k = -1$. Such a typical behaviour appears for the first time and perhaps, Hamiltonian constraint, which is a special feature of gravitation is responsible for such contradiction. The question that arises, whether $k = \pm 1$ admits any other symmetry or not. As far as Nöther symmetry is concerned we can say that the techniques we have followed to find such symmetry will not yield anything further. This is because equation (12) cannot be solved to obtain f in closed form. However, still there might exist symmetry which can be studied from the field equations only. To show this we choose a form of equations $f = -\epsilon\phi^2$, keeping ϵ arbitrary and study whether any symmetry exists for $\epsilon \neq 1/12$. Thus the action is now

$$A = \int d^4x \sqrt{-g} [-\epsilon\phi^2 R - \frac{1}{2}(\phi_{,\mu}\phi^{,\mu} - V(\phi))], \quad (34)$$

which reduces to

$$A = \int [6\epsilon a \dot{a}^2 \phi^2 + 12\epsilon a^2 \phi \dot{a} \dot{\phi} - 6\epsilon k a \phi^2 + \frac{1}{2}(a^3 \dot{\phi}^2 - a^3 V(\phi))] dt + \text{surface-term}. \quad (35)$$

Thus the field equations are

$$2\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + 4\frac{\dot{a}\dot{\phi}}{a\phi} + [2 - \frac{1}{4\epsilon}]\frac{\dot{\phi}^2}{\phi^2} + 2\frac{\ddot{\phi}}{\phi} + \frac{k}{a^2} + \frac{V}{2\epsilon\phi^2} = 0, \quad (36)$$

$$\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{\dot{a}\dot{\phi}}{4\epsilon a\phi} + \frac{\ddot{\phi}}{12\epsilon\phi} + \frac{k}{a^2} + \frac{V'}{12\epsilon\phi} = 0, \quad (37)$$

$$\frac{\dot{a}^2}{a^2} + 2\frac{\dot{a}\dot{\phi}}{a\phi} + \frac{\dot{\phi}^2}{12\epsilon\phi^2} + \frac{k}{a^2} + \frac{V(\phi)}{6\epsilon\phi^2} = 0. \quad (38)$$

One can now proceed to find Nöther symmetry for the Lagrangian (35), but then it will end up with the same result, i.e., for $k = \pm 1$, Nöther symmetry exists for $\epsilon = 1/12$, which not only makes the Lagrangian degenerate but also restricts V in such a way ($V = \lambda\phi^6$), that field equations are not satisfied. However, one can still study the possibility of existence of some other form of symmetry, in view of the the continuity equation, which is obtained by multiplying equation (37) by 2 and then subtracting it from equation (31) and finally adding it up with equation (38). The equation thus formed is the same equation (21) but for $f = -\epsilon\phi^2$ viz.,

$$(12\epsilon - 1)\left[\frac{\ddot{\phi}}{\phi} + \frac{\dot{\phi}^2}{\phi^2} + 3\frac{\dot{a}\dot{\phi}}{a\phi}\right] + \frac{\phi V' - 4V}{\phi^2} = 0. \quad (39)$$

It is clear from the above equation that if one chooses $\epsilon = 1/12$ then $\phi V' - 4V = 0$ i.e., $V = \lambda\phi^4$, which is the same old result. On the contrary if one starts with a quartic potential i.e. $V = \lambda\phi^4$ and restricts oneself to such situation that the Lagrangian does not turn out to be degenerate i.e. $\epsilon \neq 1/12$, then for any arbitrary ϵ one would obtain

$$\frac{\ddot{\phi}}{\phi} + \frac{\dot{\phi}^2}{\phi^2} + 3\frac{\dot{a}\dot{\phi}}{a\phi} = 0, \quad (40)$$

i.e.,

$$a^3\phi\dot{\phi} = \text{constant}. \quad (41)$$

Hence we obtain a constant of motion, which implies that the Lagrangian has got a symmetry for all $k = 0, \pm 1$, that does not make it degenerate. Clearly, $f = \epsilon\phi^2$ does not satisfy equation (12) for arbitrary Q , i.e., the symmetry obtained just by studying the continuity equation does not belong to Nöther class. This result implies that not all (dynamical?) symmetries of Lagrangian are revealed by studying Nöther theorem.

4 Conclusion

The two very important outcomes of this paper are as follows : (1) The Lagrangian (2) seems to admit Nöther symmetry $\mathcal{L}_x L = 0$ for ($k = \pm 1$) $f = -1/12(\phi - \phi_0)^2$ and $V = \lambda(\phi - \phi_0)^6$. However these forms of f and V do not satisfy the field equations except for very special case viz., $V = 0$ and $k = -1$ yielding nontrivial solutions of the field equations. For $f = -1/12(\phi - \phi_0)^2$, the field equations are satisfied for $V = \lambda(\phi - \phi_0)^4$, but this does not yield a conserved current for $k = \pm 1$. The degeneracy of the Lagrangian is not responsible for such contradiction, since for $k = 0$, $f = -1/12(\phi - \phi_0)^2$ and $V = \lambda(\phi - \phi_0)^4$ we have obtained a constant of motion, viz., equation (30) which does not resemble with Nöther symmetry. The constraint due to the degenerate Lagrangian in this case leads to a single equation with a pair of field variables a and ϕ , that can be solved in principle assuming a particular form of ϕ and the solution is found to be nontrivial. Such situation arises quite often in general theory of relativity and the problem is tackled by assuming equation of state or some other condition. Hence we conclude that the trivial solutions viz., $a(t) = 0$ and $\phi(t) = 0$, as obtained by Capozzeillo et al [4] is not due to the constraint imposed by degenerate Lagrangian as pointed out by them, rather due to the choice of the potential $V = \lambda\phi^6$, that does not satisfy the field equations.

Situation, like Nöther symmetry exists but field equations are not satisfied has not perhaps been encountered before. The reason for such contradiction might be due to the presence of Hamiltonian constraint equation, since we have observed that this equation in particular is not satisfied by the symmetry.

(2) Though the Lagrangian (2) does not admit Nöther symmetry $\mathcal{L}_x L = 0$, yet there exists symmetry that can be studied just from the continuity equation. It has been shown in section 3, that for $k = 0, \pm 1$; $V = \lambda\phi^4$ and $f = -\epsilon\phi^2$, $\epsilon \neq 0, \neq 1/12$ being arbitrary, there exists a constant of motion and hence a symmetry in the Lagrangian. For $\epsilon \neq 0$ and $\neq 1/12$, the Hessian determinant does not vanish and hence Lagrangian is not degenerate. This symmetry is not of Nöther class since such f and V do not satisfy equation (12). This implies that even though Nöther theorem fails to extract any symmetry from the Lagrangian, continuity equation might help to see whether symmetry exists in any other form. Hence we conclude that the Nöther theorem is not the last word on (dynamical ?) symmetry.

References

- [1] R. de Ritis et al. (1990) Phys. Rev D42, 1091
- [2] R. de Ritis et al. (1991) Phys Lett A 161, 230, M. Demianski et al, (1991) Phys Rev D44, 3136
- [3] D. La and P. J. Steinhardt, (1989) Phys. Rev. Lett 62, 376, D. La, P. J. Steinhardt and E. W. Bertschinger, (1989) Phys. Lett B 231, 231, A. D. Linde, (1990) Phys. Lett B 238, 160.
- [4] S. Capozziello and R. de Ritis. (1993) Phys. Lett A 177, 1, S. Capozziello, R. de Ritis and P. Scudellaro (1994) Phys Lett A 188, 130, S. Capozziello et al (1996) Nuovo cimento B 109, 159, S. Capozziello and R. de Ritis, (1994) Class. Quantum Grav 11, 107.
- [5] K. Sundermeyer, (1982) Constrained Dynamics, Springer-Verlag.